

INVERSE TRIGONOMETRIC AND HYPERBOLIC SUMMATION FORMULAS INVOLVING GENERALIZED FIBONACCI NUMBERS

R. S. Melham and A. G. Shannon

University of Technology, Sydney, 2007, Australia

(Submitted June 1993)

1. INTRODUCTION

Define the sequences $\{U_n\}_{n=0}^{\infty}$ and $\{V_n\}_{n=0}^{\infty}$ for all integers n by

$$U_n = pU_{n-1} + U_{n-2}, \quad U_0 = 0, \quad U_1 = 1, \quad n \geq 2, \quad (1.1)$$

$$V_n = pV_{n-1} + V_{n-2}, \quad V_0 = 2, \quad V_1 = p, \quad n \geq 2. \quad (1.2)$$

Of course, these sequences can be extended to negative subscripts by the use of (1.1) and (1.2). The Binet forms for U_n and V_n are

$$U_n = \frac{\alpha^n - \beta^n}{\alpha - \beta}, \quad (1.3)$$

and

$$V_n = \alpha^n + \beta^n, \quad (1.4)$$

where

$$\alpha = \frac{p + \sqrt{p^2 + 4}}{2}, \quad \beta = \frac{p - \sqrt{p^2 + 4}}{2}. \quad (1.5)$$

Certain specializations of the parameter p produce sequences that are of interest here and Table 1 summarizes these.

TABLE 1

p	1	2	$2x$
U_n	F_n	P_n	$P_n(x)$
V_n	L_n	Q_n	$Q_n(x)$

Here $\{F_n\}$ and $\{L_n\}$ are the Fibonacci and Lucas sequences, respectively. The sequences $\{P_n\}$ and $\{Q_n\}$ are the Pell and Pell-Lucas numbers, respectively, and appear, for example, in [3], [5], [7], [11], and [17]. The sequences $\{P_n(x)\}$ and $\{Q_n(x)\}$ are the Pell and Pell-Lucas polynomials, respectively, and have been studied, for example, in [12], [14], [15], and [16].

Hoggatt and Ruggles [9] produced some summation identities for Fibonacci and Lucas numbers involving the arctan function. Their results are of the same type as the striking result of D. H. Lehmer,

$$\sum_{i=1}^{\infty} \tan^{-1} \left(\frac{1}{F_{2i+1}} \right) = \frac{\pi}{4}, \quad (1.6)$$

to which reference is made in their above-mentioned article [9]. Mahon and Horadam [13] produced identities for Pell and Pell-Lucas polynomials leading to summation formulas for Pell and Pell-Lucas numbers similar to (1.6). For example,

$$\sum_{i=0}^{\infty} \tan^{-1} \left(\frac{2}{P_{2i+1}} \right) = \frac{\pi}{2}. \quad (1.7)$$

Here, we produce similar results involving the arctan function and terms from the sequences $\{U_n\}$ and $\{V_n\}$. Some of our results are equivalent to those obtained in [13] but most are new. We also obtain results involving the arctanh function, all of which we believe are new.

2. PRELIMINARY RESULTS

We make consistent use of the following results which appear in [1] and [6]:

$$\tan^{-1} x + \tan^{-1} y = \tan^{-1} \left(\frac{x+y}{1-xy} \right), \text{ if } xy < 1, \quad (2.1)$$

$$\tan^{-1} x - \tan^{-1} y = \tan^{-1} \left(\frac{x-y}{1+xy} \right), \text{ if } xy > -1, \quad (2.2)$$

$$\tanh^{-1} x + \tanh^{-1} y = \tanh^{-1} \left(\frac{x+y}{1+xy} \right), \quad (2.3)$$

$$\tanh^{-1} x - \tanh^{-1} y = \tanh^{-1} \left(\frac{x-y}{1-xy} \right), \quad (2.4)$$

$$\tanh^{-1} x = \frac{1}{2} \log_e \left(\frac{1+x}{1-x} \right), \quad |x| < 1, \quad (2.5)$$

$$\coth^{-1} x = \frac{1}{2} \log_e \left(\frac{x+1}{x-1} \right), \quad |x| > 1, \quad (2.6)$$

$$\tan^{-1} x = \cot^{-1} \left(\frac{1}{x} \right), \quad (2.7)$$

$$\tanh^{-1} x = \coth^{-1} \left(\frac{1}{x} \right). \quad (2.8)$$

We note from (2.7) and (2.8) that all results obtained involving arctan (arctanh) can be expressed equivalently using arccot (arccoth).

If k and n are integers, and writing

$$\Delta = (\alpha - \beta)^2 = p^2 + 4, \quad (2.9)$$

we also have the following:

$$U_n^2 - U_{n+k}U_{n-k} = (-1)^{n+k} U_k^2, \quad (2.10)$$

$$V_{n+k}V_{n-k} - V_n^2 = \Delta(-1)^{n+k} U_k^2, \quad (2.11)$$

$$U_{n+k} - U_{n-k} = \begin{cases} U_k V_n, & k \text{ even}, \\ U_n V_k, & k \text{ odd}, \end{cases} \quad (2.12)$$

$$U_{n+k} + U_{n-k} = \begin{cases} U_n V_k, & k \text{ even}, \\ U_k V_n, & k \text{ odd}, \end{cases} \quad (2.13)$$

$$V_{n+k} - V_{n-k} = \begin{cases} \Delta U_k U_n, & k \text{ even}, \\ V_k V_n, & k \text{ odd}, \end{cases} \quad (2.14)$$

$$V_{n+k} + V_{n-k} = \begin{cases} V_k V_n, & k \text{ even}, \\ \Delta U_k U_n, & k \text{ odd}. \end{cases} \quad (2.15)$$

$$U_{n+2} + U_n = V_{n+1}, \quad (2.16)$$

$$U_n U_{n+2} + (-1)^n = U_{n+1}^2. \quad (2.17)$$

Identities (2.12)-(2.15) occur as (56)-(63) in [2], and the remainder can be proved using Binet forms. Indeed, (2.10) and (2.11) resemble the famous Catalan identity for Fibonacci numbers,

$$F_n^2 - F_{n+k}F_{n-k} = (-1)^{n-k} F_k^2. \quad (2.18)$$

We assume throughout that the parameter p is real and $|p| \geq 1$. If $p \geq 1$, then $\{U_n\}_{n=2}^\infty$ and $\{V_n\}_{n=1}^\infty$ are increasing sequences. If $p \leq -1$, then $\{|U_n|\}_{n=2}^\infty$ and $\{|V_n|\}_{n=1}^\infty$ are increasing sequences and, if $n > 0$, then

$$\begin{cases} U_n < 0, & n \text{ even}, \\ U_n > 0, & n \text{ odd}, \\ V_n < 0, & n \text{ odd}, \\ V_n > 0, & n \text{ even}. \end{cases} \quad (2.19)$$

Furthermore, if $|p| \geq 1$, then using Binet forms it is seen that

$$\lim_{n \rightarrow \infty} \frac{U_{n+m}}{U_n} = \lim_{n \rightarrow \infty} \frac{V_{n+m}}{V_n} = \begin{cases} \delta^m, & m \text{ even or } p \geq 1, \\ -\delta^m, & m \text{ odd and } p \leq -1, \end{cases} \quad (2.20)$$

where

$$\delta = \frac{|p| + \sqrt{p^2 + 4}}{2}. \quad (2.21)$$

3. MAIN RESULTS

Theorem 1: If n is an integer, then

$$\tan^{-1} U_{n+2} - \tan^{-1} U_n = \tan^{-1} \left(\frac{p}{U_{n+1}} \right), \quad n \text{ even}, \quad (3.1)$$

$$\tan^{-1} \left(\frac{1}{U_n} \right) + \tan^{-1} \left(\frac{1}{U_{n+2}} \right) = \tan^{-1} \left(\frac{V_{n+1}}{U_{n+1}^2} \right), \quad n \text{ odd}, n \neq -1. \quad (3.2)$$

Proof:

$$\tan^{-1} U_{n+2} - \tan^{-1} U_n = \tan^{-1} \left(\frac{U_{n+2} - U_n}{1 + U_n U_{n+2}} \right) = \tan^{-1} \left(\frac{p}{U_{n+1}} \right),$$

where we have used (2.2), (1.1), and (2.17).

To prove (3.2), proceed similarly using (2.1), (2.16), and (2.17). \square

Now, in (3.1), replacing n by $0, 2, \dots, 2n-2$, we obtain a sum which telescopes to yield

$$\sum_{i=1}^n \tan^{-1} \left(\frac{p}{U_{2i-1}} \right) = \tan^{-1} U_{2n}. \quad (3.3)$$

Similarly, in (3.2), replacing n by $1, 3, \dots, 2n-1$ yields

$$\sum_{i=1}^n (-1)^{i-1} \tan^{-1} \left(\frac{V_{2i}}{U_{2i}^2} \right) = \frac{\pi}{4} + (-1)^{n-1} \tan^{-1} \left(\frac{1}{U_{2n+1}} \right). \quad (3.4)$$

The corresponding limiting sums are

$$\sum_{i=1}^{\infty} \tan^{-1} \left(\frac{p}{U_{2i-1}} \right) = \begin{cases} \frac{\pi}{2}, & p \geq 1, \\ -\frac{\pi}{2}, & p \leq -1, \end{cases} \quad (3.5)$$

$$\sum_{i=1}^{\infty} (-1)^{i-1} \tan^{-1} \left(\frac{V_{2i}}{U_{2i}^2} \right) = \frac{\pi}{4}. \quad (3.6)$$

We note here that (3.3) and (3.4) were essentially obtained by Mahon and Horadam [13], (3.3) in a slightly different form. When $p = 1$, (3.5) reduces essentially to Lehmer's result (1.6) stated earlier.

Theorem 2: For positive integers k and n ,

$$\tan^{-1} \left(\frac{U_n}{U_{n+k}} \right) - \tan^{-1} \left(\frac{U_{n-k}}{U_n} \right) = \begin{cases} \tan^{-1} \left(\frac{(-1)^n U_k^2}{V_k U_n^2} \right), & k \text{ even}, \\ \tan^{-1} \left(\frac{(-1)^{n-1} U_k}{U_{2n}} \right), & k \text{ odd}. \end{cases} \quad (3.7)$$

Proof: Use (2.2), (2.10), and (2.13). \square

Now, in (3.7), replacing n by $k, 2k, \dots, nk$ to form a telescoping sum yields

$$\sum_{i=1}^n \tan^{-1} \left(\frac{U_k^2}{V_k U_{ik}^2} \right) = \tan^{-1} \left(\frac{U_{nk}}{U_{(n+1)k}} \right), \quad k \text{ even} \quad (3.8)$$

$$\sum_{i=1}^n \tan^{-1} \left(\frac{(-1)^{i-1} U_k}{U_{2ik}} \right) = \tan^{-1} \left(\frac{U_{nk}}{U_{(n+1)k}} \right), \quad k \text{ odd}. \quad (3.9)$$

Using (2.20), the limiting sums are, respectively,

$$\sum_{i=1}^{\infty} \tan^{-1} \left(\frac{U_k^2}{V_k U_{ik}^2} \right) = \tan^{-1}(\delta^{-k}), \quad k \text{ even}, \quad (3.10)$$

$$\sum_{i=1}^{\infty} \tan^{-1} \left(\frac{(-1)^{i-1} U_k}{U_{2ik}} \right) = \begin{cases} \tan^{-1}(\delta^{-k}), & k \text{ odd}, p \geq 1, \\ -\tan^{-1}(\delta^{-k}), & k \text{ odd}, p \leq -1. \end{cases} \quad (3.11)$$

Theorem 3: For positive integers k and n ,

$$\tan^{-1} \left(\frac{V_{n-k}}{V_n} \right) - \tan^{-1} \left(\frac{V_n}{V_{n+k}} \right) = \begin{cases} \tan^{-1} \left(\frac{\Delta(-1)^n U_k^2}{V_k V_n^2} \right), & k \text{ even}, \\ \tan^{-1} \left(\frac{(-1)^{n-1} U_k}{U_{2n}} \right), & k \text{ odd}. \end{cases} \quad (3.12)$$

Proof: Use (2.2), (2.11), and (2.15). \square

Again in (3.12), replacing n by $k, 2k, \dots, nk$ yields

$$\sum_{i=1}^n \tan^{-1} \left(\frac{\Delta U_k^2}{V_k V_{ik}^2} \right) = \tan^{-1} \left(\frac{2}{V_k} \right) - \tan^{-1} \left(\frac{V_{nk}}{V_{(n+1)k}} \right), \quad k \text{ even}, \quad (3.13)$$

$$\sum_{i=1}^n \tan^{-1} \left(\frac{(-1)^{i-1} U_k}{U_{2ik}} \right) = \tan^{-1} \left(\frac{2}{V_k} \right) - \tan^{-1} \left(\frac{V_{nk}}{V_{(n+1)k}} \right), \quad k \text{ odd}. \quad (3.14)$$

Since the left sides of (3.9) and (3.14) are the same, we can write

$$\tan^{-1} \left(\frac{U_{nk}}{U_{(n+1)k}} \right) + \tan^{-1} \left(\frac{V_{nk}}{V_{(n+1)k}} \right) = \tan^{-1} \left(\frac{2}{V_k} \right), \quad k \text{ odd}, \quad (3.15)$$

and taking limits using (2.20) gives

$$\tan^{-1}(\delta^{-k}) = \frac{1}{2} \tan^{-1} \left(\frac{2}{|V_k|} \right), \quad k \text{ odd}. \quad (3.16)$$

The limiting sum arising from (3.13) is

$$\sum_{i=1}^{\infty} \tan^{-1} \left(\frac{\Delta U_k^2}{V_k V_{ik}^2} \right) = \tan^{-1} \left(\frac{2}{V_k} \right) - \tan^{-1}(\delta^{-k}), \quad k \text{ even.} \quad (3.17)$$

Theorem 4: If $n > 2$, then

$$\tanh^{-1} \left(\frac{1}{U_n} \right) + \tanh^{-1} \left(\frac{1}{U_{n+2}} \right) = \tanh^{-1} \left(\frac{V_{n+1}}{U_{n+1}^2} \right), \quad n \text{ even,} \quad (3.18)$$

$$\tanh^{-1} \left(\frac{1}{U_n} \right) - \tanh^{-1} \left(\frac{1}{U_{n+2}} \right) = \tanh^{-1} \left(\frac{p}{U_{n+1}} \right), \quad n \text{ odd.} \quad (3.19)$$

Proof: To prove (3.18) use (2.3), (2.16), and (2.17); (3.19) is proved similarly. \square

These results lead, respectively, to

$$\sum_{i=1}^n (-1)^{i-1} \tanh^{-1} \left(\frac{V_{2i+3}}{U_{2i+3}^2} \right) = \tanh^{-1} \left(\frac{1}{U_4} \right) + (-1)^{n-1} \tanh^{-1} \left(\frac{1}{U_{2n+4}} \right), \quad (3.20)$$

$$\sum_{i=1}^n \tanh^{-1} \left(\frac{p}{U_{2i+2}} \right) = \tanh^{-1} \left(\frac{1}{U_3} \right) - \tanh^{-1} \left(\frac{1}{U_{2n+3}} \right). \quad (3.21)$$

Note that in Theorem 4 our assumption that $n > 2$, together with our earlier assumption that $|p| \geq 1$, is necessary to ensure that the arctanh function is defined. The corresponding limiting sums are

$$\sum_{i=1}^{\infty} (-1)^{i-1} \tanh^{-1} \left(\frac{V_{2i+3}}{U_{2i+3}^2} \right) = \tanh^{-1} \left(\frac{1}{U_4} \right), \quad (3.22)$$

$$\sum_{i=1}^{\infty} \tanh^{-1} \left(\frac{p}{U_{2i+2}} \right) = \tanh^{-1} \left(\frac{1}{U_3} \right). \quad (3.23)$$

We refrain from giving proofs for the theorems that follow, since the proofs are similar to those already given.

Theorem 5: Let $n \geq k$ be positive integers where $(k, n) \neq (1, 1)$ if $p = \pm 1$. Then

$$\tanh^{-1} \left(\frac{U_n}{U_{n+k}} \right) - \tanh^{-1} \left(\frac{U_{n-k}}{U_n} \right) = \begin{cases} \tanh^{-1} \left(\frac{(-1)^n U_k}{U_{2n}} \right), & k \text{ even,} \\ \tanh^{-1} \left(\frac{(-1)^{n-1} U_k^2}{V_k U_n^2} \right), & k \text{ odd.} \end{cases} \quad (3.24)$$

This leads to

$$\sum_{i=1}^n \tanh^{-1} \left(\frac{U_k}{U_{2ik}} \right) = \tanh^{-1} \left(\frac{U_{nk}}{U_{(n+1)k}} \right), \quad k \text{ even,} \quad (3.25)$$

$$\sum_{i=1}^n \tanh^{-1} \left(\frac{(-1)^{i-1} U_k^2}{V_k U_{ik}^2} \right) = \tanh^{-1} \left(\frac{U_{nk}}{U_{(n+1)k}} \right), \quad k \text{ odd.} \quad (3.26)$$

The corresponding limiting sums are

$$\sum_{i=1}^{\infty} \tanh^{-1} \left(\frac{U_k}{U_{2ik}} \right) = \tanh^{-1}(\delta^{-k}), \quad k \text{ even,} \quad (3.27)$$

$$\sum_{i=1}^{\infty} \tanh^{-1} \left(\frac{(-1)^{i-1} U_k^2}{V_k U_{ik}^2} \right) = \begin{cases} \tanh^{-1}(\delta^{-k}), & k \text{ odd, } p \geq 1, \\ -\tanh^{-1}(\delta^{-k}), & k \text{ odd, } p \leq -1. \end{cases} \quad (3.28)$$

Theorem 6: Let $n \geq k$ be positive integers where $(k, n) \neq (1, 1)$ if $1 \leq |p| \leq 2$. Then

$$\tanh^{-1} \left(\frac{V_{n-k}}{V_n} \right) - \tanh^{-1} \left(\frac{V_n}{V_{n+k}} \right) = \begin{cases} \tanh^{-1} \left(\frac{(-1)^n U_k}{U_{2n}} \right), & k \text{ even,} \\ \tanh^{-1} \left(\frac{\Delta(-1)^{n-1} U_k^2}{V_k V_n^2} \right), & k \text{ odd.} \end{cases} \quad (3.29)$$

The resulting sums are

$$\sum_{i=1}^n \tanh^{-1} \left(\frac{U_k}{U_{2ik}} \right) = \tanh^{-1} \left(\frac{2}{V_k} \right) - \tanh^{-1} \left(\frac{V_{nk}}{V_{(n+1)k}} \right), \quad k \text{ even,} \quad (3.30)$$

$$\sum_{i=1}^n \tanh^{-1} \left(\frac{\Delta(-1)^{i-1} U_k^2}{V_k V_{ik}^2} \right) = \tanh^{-1} \left(\frac{2}{V_k} \right) - \tanh^{-1} \left(\frac{V_{nk}}{V_{(n+1)k}} \right), \quad k \text{ odd.} \quad (3.31)$$

Since the left sides of (3.25) and (3.30) are the same, we can write

$$\tanh^{-1} \left(\frac{U_{nk}}{U_{(n+1)k}} \right) + \tanh^{-1} \left(\frac{V_{nk}}{V_{(n+1)k}} \right) = \tanh^{-1} \left(\frac{2}{V_k} \right), \quad k \text{ even,} \quad (3.32)$$

and taking limits yields

$$\tanh^{-1}(\delta^{-k}) = \frac{1}{2} \tanh^{-1} \left(\frac{2}{V_k} \right), \quad k \text{ even.} \quad (3.33)$$

This should be compared with (3.16). The limiting sum arising from (3.31) is

$$\sum_{i=1}^{\infty} \tanh^{-1} \left(\frac{\Delta(-1)^{i-1} U_k^2}{V_k V_{ik}^2} \right) = \begin{cases} \tanh^{-1} \left(\frac{2}{V_k} \right) - \tanh^{-1}(\delta^{-k}), & k \text{ odd, } p \geq 1, \\ \tanh^{-1} \left(\frac{2}{V_k} \right) + \tanh^{-1}(\delta^{-k}), & k \text{ odd, } p \leq -1. \end{cases} \quad (3.34)$$

At this point we remark that Mahon and Horadam [13] obtained results similar to our Theorems 2 and 3 and derived summation formulas from them. However, in our notation, they considered only the case k odd.

4. APPLICATIONS

We now use some of our results to obtain identities for the Fibonacci and Lucas numbers. From (3.22) and (3.23), we have

$$\sum_{i=1}^{\infty} (-1)^{i-1} \tanh^{-1} \left(\frac{L_{2i+3}}{F_{2i+3}^2} \right) = \frac{1}{2} \log_e 2, \quad (4.1)$$

$$\sum_{i=1}^{\infty} \tanh^{-1} \left(\frac{1}{F_{2i+2}} \right) = \frac{1}{2} \log_e 3. \quad (4.2)$$

In terms of infinite products, these become, respectively,

$$\prod_{i=1}^{\infty} \frac{F_{2i+3}^2 + (-1)^{i-1} L_{2i+3}}{F_{2i+3}^2 + (-1)^i L_{2i+3}} = 2, \quad (4.3)$$

$$\prod_{i=1}^{\infty} \frac{F_{2i+2} + 1}{F_{2i+2} - 1} = 3. \quad (4.4)$$

In (3.28), keeping in mind the constraints in the statement of Theorem 5 and taking $k = 3$, we obtain

$$\sum_{i=1}^{\infty} (-1)^{i-1} \tanh^{-1} \left(\frac{1}{F_{3i}^2} \right) = \frac{1}{2} \log_e \phi, \quad (4.5)$$

or

$$\prod_{i=1}^{\infty} \frac{F_{3i}^2 + (-1)^{i-1}}{F_{3i}^2 + (-1)^i} = \phi, \quad (4.6)$$

where $\phi = \frac{1+\sqrt{5}}{2}$ is the Golden Ratio.

Finally, (3.34) yields, after simplifying the right side,

$$\sum_{i=1}^{\infty} (-1)^{i-1} \tanh^{-1} \left(\frac{5}{L_{3i}^2} \right) = \frac{1}{2} \log_e (3(\phi - 1)), \quad (4.7)$$

or

$$\prod_{i=1}^{\infty} \frac{L_{3i}^2 + (-1)^{i-1} 5}{L_{3i}^2 + (-1)^i 5} = 3(\phi - 1). \quad (4.8)$$

Of course, many other examples can be given by varying the parameter k .

REFERENCES

1. M. Abramowitz, & I. A. Stegun. *Handbook of Mathematical Functions*. New York: Dover, 1972.
2. G. E. Bergum & V. E. Hoggatt, Jr. "Sums and Products for Recurring Sequences." *The Fibonacci Quarterly* **13.2** (1975):115-20.
3. M. Bicknell. "A Primer on the Pell Sequence and Related Sequences." *The Fibonacci Quarterly* **13.4** (1975):345-49.

4. O. Brugia, A. Di Porto, & P. Filipponi. "On Certain Fibonacci-Type Sums." *International Journal of Mathematical Education in Science and Technology* **22** (1991):609-13.
5. E. Cohn. "Pell Number Triples." *The Fibonacci Quarterly* **10.5** (1972):403-12.
6. C. H. Edwards, Jr., & D. E. Penny. *Calculus and Analytic Geometry*. Englewood Cliffs, NJ: Prentice Hall, 1982.
7. J. Ercolano. "Matrix Generators of Pell Sequences." *The Fibonacci Quarterly* **17.1** (1979): 71-77.
8. V. E. Hoggatt, Jr., & I. D. Ruggles. "A Primer for the Fibonacci Numbers—Part IV." *The Fibonacci Quarterly* **1.4** (1963):65-71.
9. V. E. Hoggatt, Jr., & I. D. Ruggles. "A Primer for the Fibonacci Numbers—Part V." *The Fibonacci Quarterly* **2.1** (1964):59-65.
10. A. F. Horadam. "Generating Identities for Generalized Fibonacci and Lucas Triples." *The Fibonacci Quarterly* **15.4** (1977):289-93.
11. A. F. Horadam. "Pell Identities." *The Fibonacci Quarterly* **9.3** (1971):245-52, 263.
12. A. F. Horadam & Bro. J. M. Mahon. "Pell and Pell-Lucas Polynomials." *The Fibonacci Quarterly* **23.1** (1985):7-20.
13. Bro. J. M. Mahon & A. F. Horadam. "Inverse Trigonometrical Summation Formulas Involving Pell Polynomials." *The Fibonacci Quarterly* **23.4** (1985):319-24.
14. Bro. J. M. Mahon & A. F. Horadam. "Matrix and Other Summation Techniques for Pell Polynomials." *The Fibonacci Quarterly* **24.4** (1986):290-309.
15. Bro. J. M. Mahon & A. F. Horadam. "Ordinary Generating Functions for Pell Polynomials." *The Fibonacci Quarterly* **25.1** (1987):45-56.
16. Bro. J. M. Mahon & A. F. Horadam. "Pell Polynomial Matrices." *The Fibonacci Quarterly* **25.1** (1987):21-28.
17. C. Serkland. "Generating Identities for Pell Triples." *The Fibonacci Quarterly* **12.2** (1974): 121-28.

AMS Classification Numbers: 11B37, 11B39



Mark Your Calendars Now!

Mathematics Awareness Week 1995

"Mathematics and Symmetry"

April 23-29 1995

Every year, Mathematics Awareness Week celebrates the richness and relevance of mathematics and provides an excellent opportunity to convey this message through local events. During a week-long celebration from Sunday, 23 April - Saturday, 29 April 1995, the festivities will highlight **"MATHEMATICS AND SYMMETRY"**. Mark your calendars now and plan to observe Mathematics Awareness Week in your area, school, or organization. Look for further information from the Joint Policy Board for Mathematics, national sponsor of Mathematics Awareness Week, in future issues of *The Fibonacci Quarterly*.

The Fibonacci Quarterly

Official Publication of The Fibonacci Association



[Journal Home](#) | [Editorial Board](#) | [List of Issues](#)
[How to Subscribe](#) | [General Index](#) | [Fibonacci Association](#)

Volume 33

Number 1

February 1995

CONTENTS

[Cover Page](#)

Referees 2
[Full text](#)

Clark Kimberling
The Zeckendorf Array Equals the Wythoff Array 3
[Full text](#)

W. R. Spickerman, R. L. Creech, and R. N. Joyner
On the (3, F) Generalizations of the Fibonacci Sequence 9
[Full text](#)

R. S. Melham and A. G. Shannon
Some Infinite Series Summations Using Power Series Evaluated at a Matrix 13
[Full text](#)

Julia Abrahams
Varn Codes and Generalized Fibonacci Trees 21
[Full text](#)

Roger Webster
A Combinatorial Problem with a Fibonacci Solution 26
[Full text](#)

R. S. Melham and A. G. Shannon
Inverse Trigonometric Hyperbolic Summation Formulas Involving Generalized Fibonacci Numbers 32
[Full text](#)

Morris Jack DeLeon
Sequences Related to an Infinite Product Expansion for the Square Root and Cube Root Functions 41
[Full text](#)

Marjorie Bicknell-Johnson and David A. Englund
Greatest Integer Identities for Generalized Fibonacci Sequences $\{H_n\}$, Where $H_n = H_{n-1} + H_{n-2}$ 50
[Full text](#)

Wayne L. McDaniel <i>Diophantine Representation of Lucas Sequences</i> Full text	59
R. S. Melham and A. G. Shannon <i>Some Summation Identities Using Generalized Q-Matrices</i> Full text	64
Vassil S. Dimitrov and Borislav D. Donevsky <i>Faster Multiplication of Medium Large Numbers Via the Zeckendorf Representation</i> Full text	74
Paul Thomas Young <i>Quadratic Reciprocity Via Lucas Sequences</i> Full text	78
R. S. Melham and A. G. Shannon <i>A Generalization of the Catalan Identity and Some Consequences</i> Full text	82
Edited by Stanley Rabinowitz <i>Elementary Problems and Solutions</i> Full text	85
Edited by Raymond E. Whitney <i>Advanced Problems and Solutions</i> Full text	91
Back Cover	